arXiv:hep-th/9908037v1 4 Aug 1999

CHERN-SIMONS INVARIANTS OF CLOSED HYPERBOLIC 3-MANIFOLDS

A.A. BYTSENKO, A.E. GONÇALVES, AND F.L. WILLIAMS

Abstract. The Chern-Simons invariants of irreducible $U(n)-$flat connections on compact hyperbolic 3-manifolds of the form $\Gamma\backslash\mathbb{H}^3$ are derived. The explicit formula for the Chern-Simons functional is given in terms of Selberg type zeta functions related to the twisted eta invariants of Atiyah-Patodi-Singer.

PACS: 04.62.+v; 02.40.-k

1. Introduction

A gauge Chern-Simons theory is interesting both for its mathematical novelty and for its applications for certain planar condensed matter phenomena (such as the fractional quantum Hall effect) and for non-abelian gauge models in field theory. For example, the supersymmetric index can be computed in $\mathcal{N} = 1$ super Yang-Mills theory in three spacetime dimensions with a Chern-Simons interaction at level k by making a relation to a Chern-Simons theory [1].

Recently the Chern-Simons functional has been actively studied in low dimensional topology in connection with Floer homology [2], where it was used as a Morse function. This functional has also been used in topological field theory, where the new topological invariants of oriented 3-manifolds and links in it were specified [3]. The topology of manifolds can be studied with the help of the path integral approach in quantum field theory, which suggests the asymptotic behaviour of invariants including the Chern-Simons invariant, eta invariant, Redemeister torsion, and so on [4, 5, 6, 7, 8, 9, 10, 11, 12]. Under such circumstances it is important to have an explicit formula for the Chern-Simons invariant.

The Chern-Simons invariants for the case of $SU(2)$ and for some classes of 3-manifolds (including the Seifert fibered 3-manifolds) have been listed in [13, 14, 15] by cutting a 3-manifold into pieces for which

---

Date: May, 1999.

First author partially supported by a CNPq grant (Brazil), RFFI grant (Russia) No 98-02-18380-a, and by GRACENAS grant (Russia) No 6-18-1997.

the invariants can be computed. The formula for the Chern-Simons invariants of irreducible $SU(n)-$flat connection on the Seifert fibered 3-manifolds has been derived in [16]. In this paper we shall compute this invariant using the index theorem for a manifold with a closed hyperbolic boundary $X_\Gamma = \Gamma\backslash\mathbb{H}^3$.

2. The Chern-Simons invariant

In this section we briefly summarize the formalism that we shall use in this paper. Our goal is to present a formula for the $U(n)$- Chern-Simons invariant of an irreducible flat connection on a closed hyperbolic 3-manifold. This invariant is defined by the Chern-Simons functional CS which can be considered as a function on a space of connections on a trivial principal bundle over a compact oriented 3-manifold X_Γ given by

$$CS(A) = \frac{1}{8\pi^2}\int_{X_\Gamma}\mathrm{Tr}\left(A\wedge dA + \frac{2}{3}A\wedge A\wedge A\right). \tag{2.1}$$

Let X be a locally symmetric Riemannian manifold with negative sectional curvature. Its universal covering $\widetilde{X}\to X$ is a Riemannian symmetric space of rank one. The group of orientation preserving isometries $\widetilde{G}$ of $\widetilde{X}$ is a connected semisimple Lie group of real rank one and $\widetilde{X} = \widetilde{G}/\widetilde{K}$, where $\widetilde{K}$ is a maximal compact subgroup of $\widetilde{G}$. The fundamental group of X acts by covering transformations on $\widetilde{X}$ and gives rise to a discrete, co-compact subgroup $\Gamma\subset\widetilde{G}$ such that $X = \Gamma\backslash\widetilde{G}/\widetilde{K}$. If G is a linear connected finite covering of $\widetilde{G}$, the embedding $\Gamma\hookrightarrow\widetilde{G}$ lifts to an embedding $\Gamma\hookrightarrow G$. Let $K\subset G$ be a maximal compact subgroup of G, then $X_\Gamma = \Gamma\backslash G/K$ is a compact manifold.

Let $\mathrm{g} = \mathfrak{k}\oplus\mathfrak{p}$ be a Cartan decomposition of the Lie algebra g of G. Let $\mathfrak{a}\subset\mathfrak{p}$ be a one-dimensional subspace and let $J = K\bigcap G_\mathfrak{a}$ be the centralizer of $\mathfrak{a}$ in K. Fixing a positive root system of $(\mathrm{g},\mathfrak{a})$ we have an Iwasawa decomposition $\mathrm{g} = \mathfrak{k}\oplus\mathfrak{a}\oplus\mathfrak{n}$. For $G = SO(m,1)$ $(m\in\mathbb{Z}_+)$, $K = SO(m)$, and $J = SO(m-1)$. The corresponding symmetric space of non-compact type is the real hyperbolic space $\mathbb{H}^m$ of sectional curvature -1. It's compact dual space is the unit $m-$ sphere.

Let $\mathbb{P} = X_\Gamma\otimes\mathfrak{G}$ be a trivial principal bundle over X_Γ with the gauge group $\mathfrak{G} = U(n)$ and let $\mathfrak{A}_{X_\Gamma} = \Omega^1(X_\Gamma,\mathfrak{g})$ be the space of all connections on $\mathbb{P}$; this space is an affine space of 1-forms on X_Γ with values in the Lie algebra $\mathfrak{g}$ of $\mathfrak{G}$. The value of the function $CS(A)$ on the space of connections $\mathfrak{A}_{X_\Gamma}$, $A\in\mathfrak{A}_{X_\Gamma}$, at a critical point can be regarded as a topological invariant of a pair (X_Γ,χ), where χ is an orthogonal representation of the fundamental group $\pi_1(X_\Gamma)\equiv\Gamma$.

Let $\mathfrak{A}_{X_\Gamma,F} = \{A \in \mathfrak{A}_{X_\Gamma} | F_A = dA + A \wedge A = 0\}$ be the space of flat connections on $\mathbb{P}$. An important formula related to the integrand in (2.1) is

$$d\mathrm{Tr}\left(A \wedge dA + \frac{2}{3} A \wedge A \wedge A\right) = \mathrm{Tr}\left(F_A \wedge F_A\right). \tag{2.2}$$

Eq. (2.2) gives another approach to the Chern-Simons invariant. Indeed, let M be an oriented 4-manifold with boundary $\partial M = X_\Gamma$. One can extend $\mathbb{P}$ to a trivial $\mathfrak{G}-$ bundle over M; then Stokes' theorem gives

$$CS(A) = \frac{1}{8\pi^2} \int_M \mathrm{Tr}\left(F_{\widetilde{A}} \wedge F_{\widetilde{A}}\right), \tag{2.3}$$

where $\widetilde{A}$ is any extension of A over M. Eq. (2.3) can be viewed as a generalization of the Chern-Simons invariant to the case in which $\mathbb{P}$ is a non-trivial $U(n)-$ bundle over X_Γ [16]. In fact we shall show that the Chern-Simons functional is well-defined modulo$(\mathbb{Z}/2)$ in the case of a $U(n)-$ connection.

3. Invariants of closed hyperbolic manifolds

Let $\mathbb{E}_\chi = \widetilde{X} \otimes_\chi \mathbb{C}^n$ be a flat vector bundle, and let $\widetilde{\mathbb{P}}$ be a principal $U(n)-$ bundle over X_Γ which is an extension of $\mathbb{P}$. Any 1-dimensional representation χ of Γ factors through a representation of $H^1(X_\Gamma, \mathbb{Z})$. It can be shown that for a unitary representation $\chi : \Gamma \to U(n)$, the corresponding flat vector bundle $\mathbb{E}_\chi$ is topologically trivial ($\mathbb{E}_\chi \cong X_\Gamma \otimes \mathbb{C}^n$) iff $\det \chi|_{\mathrm{Tor}^1} : \mathrm{Tor}^1 \to U(1)$ is the trivial representation. Here Tor^1 is the torsion part of $H^1(X_\Gamma, \mathbb{Z})$ and $\det \chi$ is a 1-dimensional representation of Γ defined by $\det \chi(\gamma) := \det(\chi(\gamma))$, for $\gamma \in \Gamma$.

For any representation $\chi : \Gamma \to U(n)$ one can construct a vector bundle $\widetilde{\mathbb{E}}_\chi$ over a certain 4-manifold M with boundary $\partial M = X_\Gamma = \Gamma\backslash\mathbb{H}^3$ which is an extension of a flat vector bundle $\mathbb{E}_\chi$ over $\Gamma\backslash\mathbb{H}^3$. Let $\widetilde{A}_\chi$ be any extension of a flat connection A_χ corresponding to χ. The index theorem of Atiyah-Patodi-Singer for the twisted Dirac operator $D_{\widetilde{A}_\chi}$ [17, 18, 19] is given by

$$\mathrm{Ind}\left(D_{\widetilde{A}_\chi}\right) = \int_M \mathrm{ch}(\widetilde{\mathbb{E}}_\chi)\widehat{A}(M) - \frac{1}{2}(\eta(0, \mathcal{O}_\chi) + h(0, \mathcal{O}_\chi)), \tag{3.1}$$

where $h(0, \mathcal{O}_\chi)$ is the dimension of the space of harmonic spinors on X_Γ ($h(0, \mathcal{O}_\chi) = \dim\ker \mathcal{O}_\chi$ = multiplicity of the 0-eigenvalue of $\mathcal{O}_\chi$ acting on X_Γ); $\mathcal{O}_\chi$ is a Dirac operator on X_Γ acting on spinors with coefficients in χ (for details see Appendices A and B).

The Chern-Simons invariants of $\Gamma\backslash\mathbb{H}^3$ can be derived from Eq. (3.1) as follows. For a closed manifold M we have

$$-\frac{1}{8\pi^2}\mathrm{Tr}\left(F_{\widetilde{A}_\chi}\wedge F_{\widetilde{A}_\chi}\right)=\mathrm{ch}_2(\widetilde{\mathbb{E}}_\chi), \tag{3.2}$$

where $\mathrm{ch}_2(\widetilde{\mathbb{E}}_\chi)$ is the second Chern character of $\widetilde{\mathbb{E}}_\chi$, which is expressed in terms of the first and second Chern classes:

$$\mathrm{ch}_2(\widetilde{\mathbb{E}}_\chi)=\frac{1}{2}c_1(\widetilde{\mathbb{E}}_\chi)^2-c_2(\widetilde{\mathbb{E}}_\chi). \tag{3.3}$$

The Chern character and the $\widehat{A}-$ genus are given by

$$\begin{aligned}\mathrm{ch}(\widetilde{\mathbb{E}}_\chi)&=\mathrm{rank}\,\widetilde{\mathbb{E}}_\chi+c_1(\widetilde{\mathbb{E}}_\chi)+\mathrm{ch}_2(\widetilde{\mathbb{E}}_\chi)\\&=\dim\chi+c_1(\widetilde{\mathbb{E}}_\chi)+\mathrm{ch}_2(\widetilde{\mathbb{E}}_\chi),\end{aligned} \tag{3.4}$$

and

$$\widehat{A}(M)=1-\frac{1}{24}p_1(M). \tag{3.5}$$

Thus we have

$$\begin{aligned}\mathrm{ch}(\widetilde{\mathbb{E}}_\chi)\widehat{A}(M)&=\left(\dim\chi+c_1(\widetilde{\mathbb{E}}_\chi)+\mathrm{ch}_2(\widetilde{\mathbb{E}}_\chi)\right)\left(1-\frac{1}{24}p_1(M)\right)\\&=\dim\chi+c_1(\widetilde{\mathbb{E}}_\chi)+\mathrm{ch}_2(\widetilde{\mathbb{E}}_\chi)-\frac{\dim\chi}{24}p_1(M).\end{aligned} \tag{3.6}$$

The integral over the four manifold M takes the form

$$\begin{aligned}\int_M\mathrm{ch}(\widetilde{\mathbb{E}}_\chi)\widehat{A}(M)&=\int_M\mathrm{ch}_2(\widetilde{\mathbb{E}}_\chi)-\frac{\dim\chi}{24}\int_M p_1(M)\\&=-\frac{1}{8\pi^2}\int_M\mathrm{Tr}\left(F_{\widetilde{A}_\chi}\wedge F_{\widetilde{A}_\chi}\right)-\frac{\dim\chi}{24}\int_M p_1(M),\end{aligned} \tag{3.7}$$

and we have

$$\begin{aligned}\mathrm{Ind}\left(D_{\widetilde{A}_\chi}\right)&=-\frac{1}{8\pi^2}\int_M\mathrm{Tr}\left(F_{\widetilde{A}_\chi}\wedge F_{\widetilde{A}_\chi}\right)\\&-\frac{\dim\chi}{24}\int_M p_1(M)-\frac{1}{2}(\eta(0,\mathcal{O}_\chi)+h(0,\mathcal{O}_\chi)).\end{aligned} \tag{3.8}$$

For a trivial representation χ_0 one can take a trivial flat connection $\widetilde{A}\equiv\widetilde{A}_{\chi_0}$; then $F_{\widetilde{A}_{\chi_0}}=0$ and for this choice we get

$$\mathrm{Ind}\left(D_{\widetilde{A}_{\chi_0}}\right)=-\frac{1}{24}\int_M p_1(M)-\frac{1}{2}\left(\eta(0,\mathcal{O})+h(0,\mathcal{O})\right). \tag{3.9}$$

Using Eqs. (3.8) and (3.9) one can obtain

$$\mathrm{Ind}\left(D_{\widetilde{A}_\chi}\right) - \mathrm{dim}\chi\mathrm{Ind}\left(D_{\widetilde{A}_{\chi 0}}\right) = -\frac{1}{8\pi^2}\int_M \mathrm{Tr}\left(F_{\widetilde{A}_\chi} \wedge F_{\widetilde{A}_\chi}\right)$$
$$-\frac{1}{2}\left(\eta(0,\mathcal{O}_\chi) - \dim\chi\eta(0,\mathcal{O})\right) \qquad \mathrm{modulo}(\mathbb{Z}/2), \tag{3.10}$$

and

$$CS(\chi) \equiv \frac{1}{2}\left(\mathrm{dim}\chi\eta(0,\mathcal{O}) - \eta(0,\mathcal{O}_\chi)\right) \qquad \mathrm{modulo}(\mathbb{Z}/2). \tag{3.11}$$

We are now able to use the formulae (C.3) and (C.6) of Appendix C for the eta invariant in Eq. (3.11). It follows that

$$Z(0,\mathcal{O}_\chi) = Z(0,\mathcal{O})^{\dim\chi}e^{-2\pi i CS(\chi)}. \tag{3.12}$$

Finally the Chern-Simons functional takes the form

$$CS(\chi) = \frac{1}{2\pi i}\log\left[\frac{Z(0,\mathcal{O})^{\dim\chi}}{Z(0,\mathcal{O}_\chi)}\right] \qquad \mathrm{modulo}(\mathbb{Z}/2). \tag{3.13}$$

4. Concluding remarks

For an irreducible $U(n)-$ representation formula (3.13) gives the values of the Chern-Simons invariants. This is the main result of the paper. The formula for the Chern-Simons functional involves a Selberg type zeta function $Z(0,\mathcal{O}_\chi)$ associated with twisted eta invariants of Atiyah-Patodi-Singer [17, 20, 21]. In the case of a $U(n)-$ connection this is a well-defined functional, modulo($\mathbb{Z}/2$). Note also that the explicit results (3.12) and (3.13) can be very important for the study of the relation between quantum invariant for an oriented 3-manifold, defined with the help of a representation theory of quantum group [22], and Witten's invariant [3], related to the path integral approach. We hope to return to this study elsewhere.

5. Appendix A. Dirac operators and the eta invariant

We recall in this section some standard material on Dirac bundles (for details see, for example, Ref. [23]). Let M be an even-dimensional orientable Riemannian manifold with boundary $\partial M \equiv Y$ and let $\mathbb{E} \to M$ be a hermitian vector bundle equipped with a compatible connection ∇. Let $\mathbb{C}\ell(M)$ denote the complexified Clifford bundle. We suppose that there is a bundle map from $\mathbb{C}\ell(M) \to \mathrm{End}\,\mathbb{E}$ which is an algebra homomorphism on each fiber and which covers the identity:

$$\begin{tikzcd} \mathbb{C}\ell(M) \arrow[rr] \arrow[dr] & & \mathrm{End}\,\mathbb{E} \arrow[dl] \\ & M & \end{tikzcd} \qquad (A.1)$$

is commutative. Denote by $\mathbb{S}$ the spin bundles associated to the two half-spin representation of Spin(dim$M$); then $\mathbb{S} = \mathbb{S}^+ \oplus \mathbb{S}^-$ where $\mathbb{S}^\pm$ are half-spin bundles. ∇ induces a connection on $\mathbb{S}\otimes\mathbb{E}$ which is compatible with both the metric and the $\mathbb{Z}_2-$ grading. The latter connection canonically defines a Dirac operator $D : C^\infty(\mathbb{S}^+ \otimes \mathbb{E}) \rightarrow C^\infty(\mathbb{S}^- \otimes \mathbb{E})$ described by

$$D : C^\infty(\mathbb{S}^+_{\mathbb{E}}) \xrightarrow{\nabla} C^\infty(T^*M \otimes \mathbb{S}^+_{\mathbb{E}}) \xrightarrow{\mathfrak{M}} C^\infty(\mathbb{S}^-_{\mathbb{E}}), \qquad (A.2)$$

where $\mathfrak{M} : T^*M \longrightarrow \mathrm{Hom}(\mathbb{S}^+_{\mathbb{E}}, \mathbb{S}^-_{\mathbb{E}})$ denotes the Clifford multiplication and $\mathbb{S}^\pm_{\mathbb{E}} \overset{def}{=} \mathbb{S}^\pm \otimes \mathbb{E}$.

Suppose the metric on a neighborhood $\mathfrak{U} \simeq [0,1)\otimes Y$ of the boundary is a product metric. Then near the boundary Y one has a separation of variables which leads to the representation $D = \sigma\left(\frac{\partial}{\partial x} + \mathcal{O}\right)$, where the decomposition $\phi = dx \wedge \phi_1(x) + \phi_2(x)$ of a smooth form ϕ is used. Thus D acts on $C^\infty\left((0,1), \Omega^p(Y) \oplus \Omega^p(Y)\right)$, $\Omega^p(Y)$ being the space of smooth $p-$ forms on Y. A little computation shows that $\mathcal{O}$ is an elliptic self-adjoint map on $\Omega^p(Y)$ defined by $\mathcal{O}\phi = (-1)^{k+p+1}(\varepsilon * d - d*)\phi$ [17], where the symbol $*$ denotes the duality operator on Y, $k = (\dim Y)/4$ and ϕ is either a $2p-$ form ($\varepsilon = 1$) or a $(2p-1)-$ form ($\varepsilon = -1$). The operator $\mathcal{O}$ maps even (odd) forms to even (odd) forms and can therefore be decomposed as follows: $\mathcal{O} = \mathcal{O}^{ev} \oplus \mathcal{O}^{odd}$, where $\mathcal{O}^{ev}$ is isomorphic to $\mathcal{O}^{odd}$ [17].

Recall some standard material on the eta invariant of a self-adjoint elliptic differential operator acting on a compact manifold Y. For details we refer the reader to Refs. [17, 18, 19] where the eta invariant was introduced in connection with the index theorem for a manifold with boundary. To define a spectral invariant which measures the asymmetry of the spectrum Spec($\mathcal{O}$) of an operator $\mathcal{O}$, one starts with next proposition.

Proposition A.1. *The holomorphic function*

$$\eta(s,\mathcal{O}) \overset{def}{=} \sum_{\lambda\in \mathrm{Spec}\mathcal{O}\backslash\{0\}} \mathrm{sgn}(\lambda)|\lambda|^{-s} = \mathrm{Tr}\left(\mathcal{O}\left(\mathcal{O}^2\right)^{-(s+1)/2}\right), \qquad (A.3)$$

is well defined for all $\Re s \gg 0$ *and extends to a meromorphic function on* $\mathbb{C}$.

Indeed, from the asymptotic behaviour of the heat operator at $t = 0$, $\mathrm{Tr}\left(\mathcal{O}\exp\left(-t\mathcal{O}^2\right)\right) = O(t^{1/2})$ [24] and from the identity

$$\eta(s,\mathcal{O}) = \frac{1}{\Gamma\left(\frac{s+1}{2}\right)}\int_{\mathbb{R}_+} \mathrm{Tr}\left(\mathcal{O}e^{-t\mathcal{O}^2}\right) t^{(s-1)/2}dt, \qquad (A.4)$$

it follows that $\eta(s,\mathcal{O})$ admits a meromorphic extension to the whole $s-$plane, with at most simple poles at $s = (\dim Y - k)/(\mathrm{ord}\,\mathcal{O})$ $(k \in \mathbb{Z}_+)$ and locally computable residues.

It has been established that point $s = 0$ is not a pole, which makes it possible to define the eta invariant of $\mathcal{O}$ by $\eta(0,\mathcal{O})$. It also follows directly that $\eta(0,-\mathcal{O}) = -\eta(0,\mathcal{O})$ and $\eta(0,\lambda\mathcal{O}) = \eta(0,\mathcal{O})$, $\forall\lambda > 0$. One can attach the eta invariant to any operator of Dirac type on a compact Riemannian manifold of odd dimension. Dirac operators on even dimensional manifolds have symmetric spectrums and, therefore, trivial eta invariants. Because $\mathcal{O}^{ev}$ is isomorphic to $\mathcal{O}^{odd}$, we have

$$\eta(s,\mathcal{O}^{ev}) = \eta(s,\mathcal{O}^{odd}) = \frac{1}{2}\eta(s,\mathcal{O}). \qquad (A.5)$$

6. Appendix B. Closed geodesics on negative curvature manifold and Selberg type zeta function

Let Y be a compact oriented $(4m-1)-$ dimensional Riemannian manifold of constant negative curvature. A remarkable formula relating $\eta(s,B)$, $B \equiv \mathcal{O}^{ev}$, to the closed geodesics on Y has been derived by Millson [20]. More explicitly, Millson proved the following result for a Selberg type (Shintani) zeta function.

Theorem B.1 (Millson [20]). *Define a zeta function by the following series which is absolutely convergent for* $\Re\, s > 0$

$$\log Z(s,B) \overset{def}{=} \sum_{[\gamma]\neq 1} \frac{\mathrm{Tr}\tau_\gamma^+ - \mathrm{Tr}\tau_\gamma^-}{|\det(I - P_h(\gamma))|^{1/2}}\frac{e^{-s\ell(\gamma)}}{m(\gamma)}, \qquad (B.1)$$

where $[\gamma]$ *runs over the nontrivial conjugacy classes in* $\Gamma = \pi_1(Y)$, $\ell(\gamma)$ *is the length of the closed geodesic* c_γ *(with multiplicity* $m(\gamma)$*) in the free homotopy class corresponding to* $[\gamma]$, $P_h(\gamma)$ *is the restriction of the linear Poincaré map* $P(\gamma) = d\Phi_1$ *at* $(c_\gamma, \dot{c}_\gamma) \in TY$ *to the directions* *normal to the geodesic flow* Φ_t *and* $\tau_\gamma^\pm$ *is parallel translation around* c_γ *on* $\Lambda_\gamma^\pm = \pm i$ *eigenspace of* $\sigma_B(\dot{c}_\gamma)$ *(*σ_B *denoting the principal symbol of* B*). Then* $Z(s,B)$ *admits a meromorphic continuation to the entire complex plane, which in particular is holomorphic at* 0*. Moreover*

$$\log Z(0,B) = \pi i\eta(0,B). \qquad (B.2)$$

In fact $Z(s,B)$ satisfies the functional equation

$$Z(s,B)Z(-s,B) = e^{2\pi i\eta(0,B)}. \qquad (B.3)$$

7. Appendix C. Twisted eta invariants associated to locally symmetric spaces

Milson's formulae have been extended by Moscovici and Stanton [21] to Dirac type operators (acting in non-positively curved locally symmetric manifolds), even with additional coefficients in locally flat bundles. We now present the main results obtained in [21].

Let $\widetilde{Y}$ denote be a simply connected cover of Y which is a symmetric space of noncompact type, and let $\widetilde{\mathbb{E}}$ denote the pull-back to $\widetilde{Y}$ for any vector bundle $\mathbb{E}$ over Y. Bundles which will be considered satisfy a local homogeneity condition. Namely, a vector bundle $\mathbb{E}$ over Y is $\mathcal{G}-$ locally homogeneous, for some Lie group $\mathcal{G}$, if there is a smooth action of $\mathcal{G}$ on $\mathbb{E}$ which is linear on the fibers and covers the action of $\mathcal{G}$ on $\widetilde{Y}$. Standard constructions from linear algebra applied to any $\mathcal{G}-$ locally homogeneous $\mathbb{E}$ give in natural way corresponding $\mathcal{G}-$ locally homogeneous vector bundles. In particular, all bundles $TY, \mathbb{C}\ell(Y), \mathrm{End}\,\mathbb{E} \simeq \mathbb{E}^* \otimes \mathbb{E}$ are $\mathcal{G}-$ locally homogeneous [21]. We shall require also that all constructions associated with $\mathcal{G}-$ locally homogeneous bundles to be $\mathcal{G}-$ equivariant. For example, for each g in $\mathcal{G}$ the commutative diagram is

$$\begin{tikzcd}
\mathbb{C}\ell(\widetilde{Y}) \arrow[r] \arrow[d, "g"'] & \mathrm{End}\,\widetilde{\mathbb{E}} \arrow[d, "g"] \\
\mathbb{C}\ell(\widetilde{Y}) \arrow[r] & \mathrm{End}\,\widetilde{\mathbb{E}}
\end{tikzcd} \qquad (C.1)$$

Let $\mathcal{O}$ denote a generalized Dirac operator associated to a locally homogeneous bundle $\mathbb{E}$ over Y. We shall require $\mathcal{G}-$ equivariance for $\widetilde{\nabla}$ the lift of ∇ to $\widetilde{\mathbb{E}}$ and therefore the corresponding Dirac operator is then $\mathcal{G}-$ invariant.

The main results can be stated as follows.

Theorem C.1 (Moscovici and Stanton [21]). *The following function can be defined, initially for* $\Re(s^2) \gg 0$, *by the formula*

$$\log Z(s,\mathcal{O}) \overset{def}{=} \sum_{[\gamma]\in\mathcal{E}_1(\Gamma)} (-1)^q \frac{L(\gamma,\mathcal{O})}{|\det(I-P_h(\gamma))|^{1/2}} \frac{e^{-s\ell(\gamma)}}{m(\gamma)}, \qquad (C.2)$$

where $\mathcal{E}_1(\Gamma)$ *is the set of those conjugacy classes* $[\gamma]$ *for which* Y_γ *has the property that the Euclidean de Rham factor of* $\widetilde{Y}_\gamma$ *is 1-dimensional, the number* q *is one-half the dimension of the fiber of the center bundle* $C(TY)$ *over* Y_γ, *and* $L(\gamma,\mathcal{O})$ *is the Lefschetz number (see Ref.* [25]*). Furthermore* $\log Z(s,\mathcal{O})$ *has a meromorphic continuation to* $\mathbb{C}$ *given by the identity*

$$\log Z(s,\mathcal{O}) = \log\det{}' \left(\frac{\mathcal{O}-is}{\mathcal{O}+is}\right) + \pi i\eta(s,\mathcal{O}), \qquad (C.3)$$

where $s \in i\mathrm{Spec}'(\mathcal{O}) \quad (\mathrm{Spec}(\mathcal{O})-\{0\})$, *and* $Z(s,\mathcal{O})$ *satisfies the functional equation*

$$Z(s,\mathcal{O})Z(-s,\mathcal{O}) = e^{2\pi i\eta(s,\mathcal{O})}. \qquad (C.4)$$

Suppose now $\chi : \Gamma \to U(F)$ be a unitary representation of Γ on F. The Hermitian vector bundle $\mathbb{F} = \widetilde{Y} \times_\Gamma F$ over Y inherits a flat connection from the trivial connection on $\widetilde{Y} \times F$. We specialize to the case of locally homogeneous Dirac operators $\mathcal{O} : C^\infty(Y,\mathbb{E}) \to C^\infty(Y,\mathbb{E})$ in order to construct a generalized operator $\mathcal{O}_\chi$, acting on spinors with coefficients in χ. If $\mathcal{O} : C^\infty(Y,V) \to C^\infty(Y,V)$ is a differential operator acting on the sections of the vector bundle V, then $\mathcal{O}$ extends canonically to a differential operator $\mathcal{O}_\chi : C^\infty(Y, V\otimes\mathbb{F}) \to C^\infty(Y, V\otimes\mathbb{F})$, uniquely characterized by the property that $\mathcal{O}_\chi$ is locally isomorphic to $\mathcal{O}\otimes ... \otimes\mathcal{O}$ ($\dim F$ times) [21].

One can repeat the arguments of the previous sections to construct a twisted zeta function $Z(s,\mathcal{O}_\chi)$.

Theorem C.2 (Moscovici and Stanton [21]). *There exists a zeta function* $Z(s,\mathcal{O}_\chi)$, *meromorphic on* $\mathbb{C}$, *given for* $\Re(s^2) \gg 0$ *by the formula*

$$\log Z(s,\mathcal{O}_\chi) \overset{def}{=} \sum_{[\gamma]\in\mathcal{E}_1(\Gamma)} (-1)^q \mathrm{Tr}\chi(\gamma) \frac{L(\gamma,\mathcal{O})}{|\det(I-P_h(\gamma))|^{1/2}} \frac{e^{-s\ell(\gamma)}}{m(\gamma)}; \quad (C.5)$$

moreover one has

$$\log Z(0, \mathcal{O}_\chi) = \pi i \eta(0, \mathcal{O}_\chi). \qquad (C.6)$$

References

[1] E. Witten, *"Supersymmetric Index of Three-Dimensional Gauge Theory"*, hep-th/9903005.
[2] A. Floer, Commun. Math. Phys. **118** (1988) 215.
[3] E. Witten, Commun. Math. Phys. **121** (1989) 351.
[4] G. Moore and N. Seiberg, Commun. Math. Phys. **123** (1989) 77.
[5] R. Dijkgraaf and E. Witten, Commun. Math. Phys. **129** (1990) 393.
[6] R. Kirby and P. Melvin, Invent. Math. **105** (1991) 473.
[7] D.S. Freed and R.E. Gompf, Commun. Math. Phys. **141** (1991) 79.
[8] L.S. Jeffrey, Commun. Math. Phys. **147** (1992) 563.
[9] L. Rozansky, Commun. Math. Phys. **175** (1996) 275.
[10] D.H. Adams and S. Sen, *"Partition Function of a Quadratic Functional and Semiclassical Approximation for Witten's 3-Manifold Invariant"*, hep-th/9503095.
[11] A.A. Bytsenko, L. Vanzo and S. Zerbini, Nucl. Phys. B **505** (1997) 641.
[12] A.A. Bytsenko, A.E. Gonçalves and W. da Cruz, Mod. Phys. Lett. A **13** (1998) 2453.
[13] P. Kirk and E. Klassen, Math. Ann. **287** (1990) 343.
[14] P. Kirk and E. Klassen, Commun. Math. Phys. **153** (1993) 521.
[15] D. Auckly, Math. Proc. Camb. Phil. Soc. **115** (1994) 229.
[16] H. Nishi, Inter. J. Math. **9** (1998) 295.
[17] M.F. Atiyah, V.K. Patodi and I.M. Singer, Math. Proc. Camb. Phil. Soc. **77** (1975) 43.
[18] M.F. Atiyah, V.K. Patodi and I.M. Singer, Math. Proc. Camb. Phil. Soc. **78** (1975) 405.
[19] M.F. Atiyah, V.K. Patodi and I.M. Singer, Math. Proc. Camb. Phil. Soc. **79** (1976) 71.
[20] J.J. Millson, Ann. Math. **108** (1978) 1.
[21] H. Moscovici and R.J. Stanton, Invent. Math. **95** (1989) 629.
[22] N. Reshetikhin and V.G. Turaev, Invent. Math. **103** (1991) 547.
[23] H.B. Lawson and M.L. Michelson, *"Spin Geometry"*, Princeton Mathematical Series **38**, Princeton Univ. Press, Princeton (1989).
[24] J-M. Bismut and D.S. Freed, Comm. Math. Phys. **107** (1986) 103.
[25] R. Hotta and R. Parthasarathy, Osaka J. Math. **10** (1973) 211.

Departamento de Fisica, Universidade Estadual de Londrina, Caixa Postal 6001, Londrina-Parana, Brazil; on leave from Sankt-Petersburg State Technical University, Russia *E-mail address:* abyts@fisica.uel.br

Departamento de Fisica, Universidade Estadual de Londrina, Caixa Postal 6001, Londrina-Parana, Brazil *E-mail address:* goncalve@fisica.uel.br

Department of Mathematics, University of Massachusetts, Amherst, Massachusetts 01003, USA *E-mail address:* williams@math.umass.edu